

CODING FORMS FOR SRC INDEXING

Microfiche No.	OTS05J9521		
New Doc ID	86980000186	Old Doc ID	
Date Produced	09/01/98	Date Received	10/07/98
		TSCA Section	8D
Submitting Organization	INTL ISOCYANATE INST INC		
Contractor			
Document Title	IN VIVO EVALUATION OF MDI SKIN DECONTAMINATION PROCEDURES (MANUSCRIPT OF POSTER PRESENTATION), WITH COVER LETTER DATED 9/23/1998		
Chemical Category	BENZENE 1,1'-METHYLENEBIS (ISOCYANATO- (26447-40-5)		

OFFICE OF TOXIC SUBSTANCES
CODING FORM FOR GLOBAL INDEXING

REV. 7/27/82

Microfiche No. (7) •	1	No. of Pages	2
Doc I.D.	3	Old Doc I.D.	4
Case No.(s)	5		
Date Produced (6)	6	Date Rec'd (6)	7
		Conf. Code •	8
Check One: <input type="checkbox"/> Publication <input type="checkbox"/> Internally Generated <input type="checkbox"/> Externally Generated			
Pub/Journal Name			9
			9
Author(s)			10
Organ. Name			11
Dept/Div			12
P.O. Box	13	Street No./Name	14
City	15	State	16
		Zip	17
		Country	18
MID No. (7)	19	D & B NO. (11)	20
Contractor			21
Doc Type			22
Doc Title			23
Chemical Name (300 per name)	25	CAS No. (10)	24

A 03



MR 16531

INTERNATIONAL ISOCYANATE INSTITUTE, INC.

201 Main Street, Suite 403 • La Crosse, WI 54601 • 608/796-0880 • FAX 608/796-0882

September 23, 1998

TSCA Document Processing Center (TS-790)
Office of Pollution & Toxics
Environmental Protection Agency
401 M Street, SW
201 East Tower
Washington, DC 20460

Contains No CBI

Attn: 8(d) HEALTH & SAFETY STUDY REPORTING RULE
(REPORTING)

Dear Sir or Madam:

We herewith submit a copy of a manuscript of a poster presentation that is being made at Polyurethanes Expo '98. The enclosed report does not contain any Confidential Business Information.

"In Vivo Evaluation of MDI Skin Decontamination Procedures"

by Ronald C. Wester, Xiaoying Hui, Timothy D. Landry and Howard I. Maibach

Name of Chemical Substance:	benzene 1, 1' - methylenebis (isocyanato-
Chemical Abstracts Service Number:	26447-40-5
Abbreviation:	MDI

This study was sponsored by the International Isocyanate Institute on behalf of the following:

The Dow Chemical Company
Lyondell Chemical Company
Bayer Corporation
BASF Corporation
ICI Americas, Inc.

Very truly yours,

M.J. Blankenship
Managing Director

RECEIVED
OPPT NCIC
98 OCT -7 AM 10:14

RECEIVED
OPPT CBIC
98 SEP 29 AM 11:19



86980000186

FOR POLYURETHANES EXPOS '98
9/1/98 T.D. Landry

In Vivo Evaluation of MDI Skin Decontamination Procedures

Ronald C. Wester¹, Xiaoying Hui¹, Timothy D. Landry²,
Howard I. Maibach¹

¹Department of Dermatology, UCSF

²Health and Environmental Research Laboratory,
The Dow Chemical Co.

Sponsored by The International Isocyanate Institute, Inc.

ABSTRACT

Water or soap-and-water cleaning has been a traditional first aid procedure for skin decontamination in the workplace. Methylene diphenyl diisocyanate (MDI) skin exposure is a concern because it may cause skin irritation, sensitization and may play a role in respiratory sensitization. Because MDI has higher miscibility with oil or polyglycols, there may be more effective means for decontaminating skin. To assess this, ¹⁴C-MDI was applied to one cm² sites on the abdomen of monkeys, then a cleaning procedure utilizing a cotton swab with water, soap-and-water (5% or 50%), corn oil, polypropylene glycol, or a polyglycol-based skin cleanser was performed. Following cleaning at 5 minutes, 1 hour, 4 hours, or 8 hours, cellophane tape was used to check the skin for remaining radioactivity. Although all the cleaning procedures helped remove MDI to some extent, water or soap-and-water was generally less effective than corn oil, polypropylene glycol, or polyglycol-based skin cleanser. Cleaning soon after exposure was important; water or soap-and-water in particular was less effective at 4 or 8 hours. The lower amount of MDI in the outer epidermis (stratum corneum) reflected the greater effectiveness of corn oil and suitable polyglycol materials to clean skin. Although cleaning is useful, it did not completely remove skin contamination. Good work practices, engineering controls, and personal protective equipment should be the first defense to avoid skin exposure. This study showed that a rapid response is important if skin contamination occurs and these non-traditional cleaning materials may remove MDI more effectively than soap-and-water.

Introduction

- MDI may cause skin irritation and sensitization, and may play a role in respiratory sensitization
- A very small proportion of MDI applied to skin (of rats) was absorbed into the bloodstream^a
- What is the best first aid procedure for decontaminating skin of uncured MDI?

^a Hofmann, et al., (1998). Dermal pharmacokinetics of 4,4'-MDI in rats. Report to the International Isocyanate Institute, Inc. (in preparation).

Introduction (continued)

- Harsh solvents may enhance dermal penetration
- MDI has greater miscibility in oil and polyglycol than water
- Food oil or suitable polyglycols may be more effective decontaminants than soap and water

Materials and Methods

- The rhesus monkey is a suitable model for assessing dermal absorption^b
- Utilized a grid with 24 separate one cm² sites on the abdomen of each of four monkeys
- ¹⁴C-MDI added to polymeric MDI
- 2 µl of polymeric MDI (0.5 µCi) per site
- Site cleaned at 5 min, 1 hr, 4 hr, or 8 hours

^bWester and Maibach, (1997). Comprehensive Toxicology, Elsevier.

5


Methods (continued)

- Cotton swab with test cleaner (5 times per site)
 - water
 - soap-and-water (5% or 50%)
 - polypropylene glycol (m.w. 700)
 - polyglycol-based cleanser (D-TAM^c cleanser)
 - corn oil
- The type and molecular weight of polyglycol may be important:
 - low m.w. materials might be absorbed
 - high m.w. materials may be solid, not liquid
- Tape stripping of epidermis (10 times per site)


^c CLI Laboratories

6

Results

14-C in Skin Washes

7

14-C in Outer Epidermis (tape)

8

Conclusions

- Each of the cleaning procedures helped to remove some, but not all MDI
Work practices, engineering controls, and personal protective equipment should be the first defense against skin contamination
- Washing sooner is generally more effective

9

Conclusions

- Corn oil, polypropylene glycol, and polyglycol-based cleaner are generally more effective (especially at 4 & 8 hrs) than water
More 14-C was in wash, less on epithelial surface
- Results are consistent with the relative miscibility of MDI in oil and polyglycol

10

CERTIFICATE OF AUTHENTICITY

THIS IS TO CERTIFY that the microimages appearing on this microfilm are accurate and complete reproductions of the records of U.S. Environmental Protection Agency documents as delivered in the regular course of business for microfilming.

Data produced 09 17 1999 Mary Imbeck
(Month) (Day) (Year) Camera Operator

Place Syracuse New York
(City) (State)

